




Contents lists available at ScienceDirect

Pregnancy Hypertension: An International Journal of Women's Cardiovascular Health

journal homepage: www.elsevier.com/locate/preghy

Full Length Article

Cardiovascular risk management after hypertensive disorders and diabetes during pregnancy, in a multi-ethnic population: A qualitative study among women and healthcare providers

Annemarie Y.A.M. Reilingh^{a,b,c,d,*}, Renée J. Burger^{a,c,e}, Souraya el Bachiri^a, Stephen McCarthy^f, Sanne J. Gordijn^c, Wessel Ganzevoort^{a,c}, Irene G.M. Valkengoed van^{b,d},
On behalf of on behalf of the project group: 'Obstetric history and cardiovascular health across ethnicity, in The Netherlands'

^a Amsterdam UMC Location University of Amsterdam, Department of Obstetrics and Gynecology, Meibergdreef 9, Amsterdam, the Netherlands

^b Amsterdam UMC Location University of Amsterdam, Department of Public and Occupational Health, Van der Boechorststraat 7, Amsterdam, the Netherlands

^c Amsterdam Reproduction and Development, Pregnancy and Birth, Amsterdam, the Netherlands

^d Amsterdam Public Health, Health Behaviors & Chronic Diseases, Amsterdam, the Netherlands

^e University Medical Center Groningen, University of Groningen, Department of Obstetrics and Gynecology, Hanzeplein 1, Groningen, the Netherlands

^f University College Cork, Health Information Systems Research Center, College Road, Cork T12 K8AF, Ireland

ARTICLE INFO

Keywords:

Cardiovascular disease
Ethnicity
Hypertensive disorders of pregnancy
Gestational diabetes
Patient journey map
Qualitative research

ABSTRACT

Background: Pregnancy complications like gestational diabetes and hypertensive disorders increase maternal cardiovascular risk. However, evidence on how to best implement cardiovascular risk management (CVRM) in multi-ethnic contexts remains limited. Existing studies primarily focus on white populations, despite disparities in CVD risk and the risk of pregnancy complications across ethnic groups.

Objective: This study explores experiences, barriers, and improvements in postpartum CVRM from women's and healthcare providers' perspectives, aiming to enhance cardiovascular disease prevention in multi-ethnic contexts.

Study Design: We conducted semi-structured interviews with 25 postpartum women who experienced hypertensive disorders of pregnancy or gestational diabetes, recruited to reflect diverse ethnic and socioeconomic backgrounds. Using a narrative approach, interviews covered CVRM experiences and preferences. Themes were triangulated with findings from 16 healthcare providers, including general practitioners, midwives, and specialists, discussing care delivery, optimal practices, and multi-ethnic considerations. Results were integrated in a patient journey map.

Results: Significant dropout occurred at the transition from obstetric to primary care and during long-term monitoring, especially among ethnic minorities and women with lower socioeconomic status and disease severity. Women often lacked risk awareness and missed follow-ups when self-scheduling was required. Most supported tailored cardiovascular risk education, lifestyle interventions, and proactive outreach. Healthcare providers emphasized the need for interdisciplinary communication, regional protocols, and clearer guidelines, noting variability in general practitioners' support for routine monitoring.

Conclusion: Postpartum CVRM in multi-ethnic contexts could be improved with active outreach, better follow-up utilization, culturally tailored interventions, and regional multidisciplinary protocols to streamline care and address guideline inconsistencies.

1. Introduction

Hypertensive disorders of pregnancy (HDP) and gestational diabetes

mellitus (GDM) are associated with a 2 to 4-fold increased risk of future maternal cardiovascular disease (CVD) compared to healthy pregnancy [1–3]. Since complicated pregnancies occur early during the life course,

* Corresponding author at: Van der Boechorststraat 7, Amsterdam, the Netherlands.

E-mail address: a.y.a.m.reilingh@amsterdamumc.nl (A.Y.A.M. Reilingh).

<https://doi.org/10.1016/j.preghy.2025.101203>

Received 15 November 2024; Received in revised form 27 January 2025; Accepted 9 February 2025

Available online 21 February 2025

2210-7789/© 2025 The Author(s). Published by Elsevier B.V. on behalf of International Society for the Study of Hypertension in Pregnancy. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

they provide opportunities for early risk profiling and prevention of CVD, which may contribute to lowering the burden of CVD later in life [4]. Although the increased CVD risk after HDP/GDM is well established, evidence is limited on how to best implement measures to manage and reduce the future CVD risk in women after HDP/GDM [5].

Despite the existence of national and international guidelines recommending structured postpartum cardiovascular risk management (CVRM) after HDP/GDM [6,7], previous studies have shown that implementation varies, both in healthcare provider (HCP) initiated offering and in the uptake of postpartum-CVRM [8–10]. Women's and HCPs' perspectives on postpartum-CVRM have previously been studied to improve care [11–16]. However, study participants have been predominantly of white European origin, while disparities exist between women from different ethnic groups. Women from some ethnic minority groups have higher rates of pregnancy complications [17], less favorable CVD risk profiles at younger ages and worse prognoses compared to majority populations in high-income countries [18,19]. Moreover, the studies often lacked diversity in socioeconomic background and often selected a group with high interest in the topic [12–14].

To provide adequate and targeted postpartum-CVRM to all – and potentially reduce the burden of CVD in a broader group of women – perspectives on postpartum-CVRM in a multi-ethnic context should be considered. Therefore, the objective of this study was to uncover experiences, barriers, and potential improvements, from both women's and HCPs' perspectives, regarding postpartum-CVRM in a diverse multi-ethnic context.

2. Materials and methods

We performed a qualitative study among women who experienced HDP/GDM (who we proceed to call women for conciseness) and among HCPs generally involved in postpartum care. Integrated Patient Journey Mapping was used as a framework to collect the data and organize the findings from thematic analyses [20–22]. A comprehensive description of the methods following the standard reporting of qualitative research (SQSR) guidelines is found in Supplementary Table 1 [23].

In short, we recruited women with diverse, self-reported ethnic backgrounds living in the Netherlands. Women were eligible if they were between three months to five years postpartum and had experienced HDP/GDM, based on self-report. HDP included gestational hypertension, (pre)eclampsia (PE) and Hemolysis, Elevated Liver enzymes and Low platelets (HELLP) syndrome. Women with pre-existing cardiac conditions and/or hypertension and/or type II diabetes mellitus were excluded. Findings were discussed in, and triangulated with data from, interviews with HCPs (e.g., midwives, general practitioners (GPs), GP practice nurses, obstetricians, and internal medicine specialists). We used purposive sampling to reach quota per subgroup of women (six to eight participants per group) and HCPs (three participants per group). The study was approved by the Medical Ethical Review Board of the Amsterdam University Medical Center. All the participants provided informed consent.

Semi-structured interviews with women and HCPs lasted 60 and 40 min, respectively, and were held between January 2023 and March 2024 through videoconferencing or on a location of the participant's choice. Supplementary Table 2 contains the translated topic lists. Audio and video recordings were transcribed, de-identified, quality checked and locked before analysis. Women and HCP interviews were thematically analyzed separately. Coding was done by multiple coders, mostly deductively based on the topic list while allowing for inductive coding. Subsequently, we analyzed patterns in the women's data by migration background, socioeconomic status (SES), obstetric history and disease severity. Emerging themes and subthemes from thematic analyses were discussed among the entire research team in monthly meetings and final selection was based on consensus.

3. Results

In total n = 41 participants were included. The n = 25 patient participants, consisted of n = 7 women with a Dutch background and n = 18 women with an ethnic minority background (i.e., Moroccan, African Surinamese, South Asian Surinamese, Eritrean, Somali, Ghanaian, Aruban and Turkish backgrounds; Table 1). Of the women with an ethnic minority background, n = 16 were born and raised in the Netherlands, while n = 2 were born abroad. All women, except one with an ethnic minority background, were sufficiently proficient in Dutch. The distribution of educational level and employment status differed between participants with a Dutch background and those with an ethnic minority background. Participants with a Dutch background were more often highly educated and actively employed, while participants with an ethnic minority background were more often unemployed or had a lower educational level (Table 1). The women with a Dutch background were more often parous compared to those from an ethnic minority background. The n = 16 HCP participants consisted of n = 3 midwives, GPs, GP assistants and internal medicine specialists and n = 4 obstetricians. The mean age for women and HCPs was 33.3(4.9) and 44.9 (13.2) years, respectively. Of HCPs, n = 2 were male.

In the next paragraphs, we present general findings that were similar for the total group of women and findings for HCPs on: A) Experiences with the current postpartum-CVRM care pathway and B) Preferences for CVD risk education, lifestyle interventions and routine GP monitoring. We discuss findings that were specific to women from an ethnic minority background separately in paragraph C) Focus points for postpartum-CVRM in a multi-ethnic population. Quotes from the interviews support our principal findings (Table 2). A full overview of all themes and subthemes can be found in Supplementary Table 3.

3.1. Experiences with current postpartum-CVRM care pathway

We identified four main phases in the patient journey (Fig. 1), based on women's and HCPs' descriptions combined with the organization of obstetric care and CVRM within the Dutch healthcare system: 1) The first ten days postpartum, concluded by a community midwife (home) visit. 2) One to six weeks postpartum, ending with a week 6-appointment at the community midwife or obstetrician. 3) Six weeks to 12

Table 1
Characteristics of patient sample.

	Dutch BACKGROUND n = 7	Ethnic Minority BACKGROUND n = 18
Age – Mean (Sd)	36.7 (0.9)	32 (4.8)
Socioeconomic Status		
Educational Level High – N (%)	5 (71.4)	8 (44.4)
Educational Level Low to Intermediate – N (%)	1 (14.3)	10 (55.5)
Paid Work – N (%)	7 (100)	10 (55.5)
Parity (Nulliparous) – N (%)	2 (28.6)	14 (77.8)
Self-Reported Diagnosis**		
Gestational Hypertension – N (%)	1 (14.3)	1 (5.5)
(Pre)Eclampsia – N (%)	4 (28.6)	14 (66.6)
– of which HELLP Syndrome – N (%)	2 (28.6)	2 (11.1)
Gestational Diabetes Mellitus – N (%)	2 (28.6)	11 (61.1)
Pregnancy duration (weeks) – median (range)	37 (29–40)	37 (23–42)
Cesarean section – N (%)	4 (57.1)	5 (27.7)
time between birth and interview (months) – median (range)	11 (5–48)	16 (3–60)

Abbreviations: HELLP = Hemolysis, Elevated Liver enzymes and Low Platelets.
*Parity at time of index pregnancy.
**Numbers do not add up, self-reported hypertensive disorders of pregnancy, combined with & gestational diabetes mellitus are included.

Table 2

Quotes from patients and providers.

PATIENT AND PROVIDER QUOTES	
1.	<i>"And I think that's a strange thing, too, because I was not all there. I wasn't myself [...] I would say no, I missed [CVD risk information]..Because I'm now a member of the HELLP foundation. That's how I found out. Because I was so sick. Even after [HELLP], I stayed weaker. Maybe they told me, but I wasn't all there."</i> ~ Participant with acute HELLP syndrome, ethnic minority background
2.	<i>"No, I didn't have a follow-up check. That's actually also because she said you have to schedule [the appointment] yourself, but I didn't really feel like it. So, in the end it didn't happen."</i> ~ Participant with GDM, ethnic minority background
3.	<i>"You know that people who had a hypertensive disorder during pregnancy that their [cardiovascular diseases] risk is increased later on. But what hasn't been shown, or what we don't know is what is sensible care after, right? How often you have to check them to see if it's okay? To medicate so young or not? There's just no evidence for that. So that also makes it difficult to offer structural care for it."</i> ~ GP
4.	<i>"If someone had an otherwise uncomplicated delivery and the blood pressure was all normal, they went home without medication, then from the hospital they won't get the call for the follow-up check, by default [...] There are also [midwife] practices that, only if they themselves have accompanied the delivery, they see those people for follow-up checks. So, I suspect that those people who have indeed been in the hospital, but then went home fairly OK, or just without complications, that those are people who get lost between the cracks."</i> ~ Community midwife
5.	<i>"I had wanted to hear it [cardiovascular diseases risk information], so I know what to expect. Because now that I know, I know what to expect and I can prepare myself. I can start living healthier, I can do all kinds of things."</i> ~ Participant with PE, ethnic minority background
6.	<i>"No, I really prefer someone talking to me and to explain clearly. Because nine times out of ten I put a leaflet in my bag and never read it again."</i> ~ Participant with GDM/PE/HELLP, ethnic minority background
7.	<i>"...when they are 34, 35 weeks pregnant, so to speak, and have severe preeclampsia, yes, there is already quite a lot coming at them. [...] what I particularly go into at that moment is what is going on right now and how we are going to treat it. And the long-term [cardiovascular diseases risk] consequences, I think that's more for when they come for the follow-up visit [...]"</i> ~ Obstetrician
8.	<i>"Well, everybody knows that. Everybody knows that you shouldn't smoke, you shouldn't drink, you should exercise enough, you should eat healthily. [...] You know, it's just such standard textbook stuff. [...] People need more tools than just the standard lines."</i> ~ Participant with acute HELLP, Dutch background
9.	<i>"I sincerely believe that with the current care burden, how society is developing, with an aging population and more chronic diseases and obesity, we need to have a discussion because it will soon be impossible to sustain. Because there just won't be enough care providers left in institutions to do all those jobs. [...] So just to be clear, yes, we as healthcare providers absolutely have a responsibility and the task there. We need to look at how we can manage it within the healthcare system as good as possible, so that every woman who is entitled to it receives good aftercare and also is detected in a timely manner if they cross the margins. But I also think there is a task and a duty for the individual and for society."</i> ~ Internal medicine specialist
10.	<i>"Because she [the GP] had said to me: 'You have to have your blood pressure measured once a year but, you have to make an appointment yourself'. Then, yeah, if you are busy with kids and work, you think: 'I'll see'. But if someone calls me and says it's time to check my blood pressure, then, yes, that's much easier [...] no, it's not that I think like: 'oh, I have to make an appointment next week because I have to measure my blood pressure.' If I have no symptoms, then I assume that I am doing just fine".</i> ~ Participant with GDM/PE, ethnic minority background
11.	<i>"I'm very aware of [recommendations for cardiovascular diseases risk monitoring], but I can imagine that women who are either low-literate or don't speak the language or have some other barrier or obstacle in their language skills or something like that. That those are less engaged, because they may not quite understand, either what to do or when to come back for another appointment. [...] You have to figure it out yourself and you don't know the infrastructure, for example. You haven't been in the Netherlands that long yet. You can imagine that those women have a worse outcome because they don't know when they have to have blood tests again."</i> ~ Participant with GDM, ethnic minority background
12.	<i>"Well, I know that people of Moroccan descent eat a lot of bread. If someone develops diabetes [...] I do explain that bread contains a lot of carbs. I do try to make them aware. [...] the Hindustani have a lot of white rice, you know. So that's kind of hard [...] I wonder if they manage to adjust."</i> ~ GP practice nurse
13.	<i>"Certainly, the people who are less fluent in Dutch are not going to call the doctor themselves to make an appointment. [...] They might barely understand what the problem is. [...] but then to take that step to call the doctor yourself, calling the doctor at all is often complicated. [...] Well, I think, to put it very bluntly: The highly educated Dutch women who are well informed and really understand what the problem is, that they can cope with it. But the people who are less fluent in Dutch, or less educated, who just don't see the problem at all, will never take that step. While in terms of lifestyle, these are also the people who are often less healthy and more at risk."</i> ~ Community midwife

Abbreviations: GDM = gestational diabetes mellitus, GP = general practitioner, PE = preeclampsia, HELLP = Hemolysis, Elevated Liver enzymes and Low Platelets,

months postpartum, during which people may receive care at the GP, when indicated. 4) Beyond one year postpartum, routine monitoring by GP, annually in the first five years postpartum followed by once every three years, according to clinical guidelines [5].

Many women did not recall receiving CVD risk education in any phase of the care pathway with some heterogeneity depending on ethnic background. Most women received lifestyle advice, varying from general advice to in-depth personal counselling. While lifestyle advice was often already initiated during pregnancy in women with GDM, many did not follow through postpartum. Less than half of eligible women (those past one year postpartum) received routine monitoring by their GP. Compared to women with GDM, those with HDP were less often informed and less aware of the recommendations for lifestyle improvements and GP monitoring. Here, we also saw a different pattern from women from the majority compared to minority backgrounds. Moreover, women with HDP reported that the disease itself had affected information retention, motivation and initiative concerning CVRM (e.g., through impacted energy levels, mental health, work participation and memory; Table 2, quote 1).

The patient journey map shows the variation in received/provided CVD risk education, lifestyle advice and (routine) monitoring, and barriers and solutions along the care pathway (Fig. 1). The lack of received care may partly relate to the reported dropout during the care pathway, especially at phase 2 (week 6-appointment) and phase 4 (routine GP monitoring). Those missing the week 6-appointment often had an ethnic minority background and/or had lower disease severity (Table 2, quote 2). Women indicated that they were unaware about the potential important health information provided during these visits, the

fact that they had to make appointments themselves, or that they were unsure about where to receive care after transfers between hospitals. Negative experiences during pregnancy and birth, like the perception of not been taken seriously or the lack of shared decision making, sometimes led to care avoidance, including for postpartum-CVRM.

HCPs confirmed patient findings on CVD risk education, lifestyle intervention and GP monitoring. All HCPs indicated having provided CVD risk education and lifestyle interventions to women after GDM and/or HDP, but to varying degrees. They were more likely to do so after GDM than HDP, partly because of implemented guidelines on diabetes prevention, which include advice on CVRM after GDM. In general, most HCPs see benefits of early prevention for young women but acknowledge the burden on the healthcare system as a barrier. Some GPs were concerned about fragmented CVRM guidelines for specific patient groups and feel there is insufficient evidence for monitoring and lifestyle interventions after HDP/GDM (Table 2, quote 3). Many HCPs also recognized the frequent dropouts and confirm that the reliance on women's own initiative to make appointments may be a contributor, especially for women with a minority background. Some HCPs implemented active outreach with repeated calls, home visits and active scheduling of appointments. However, limited time and financial compensation prevent universal adoption of these strategies.

System barriers may also be a potential contributor. For instance, most HCPs assumed women have a week 6-appointment with either an obstetrician or with a community midwife. However, midwives indicated that this is not standard practice, potentially leading to women experiencing a gap in care (Table 2, quote 4). Additionally, GPs highlighted that while a recent multidisciplinary guideline calls for follow-up




Fig. 1. Patient Journey Map Postpartum Cardiovascular Risk Management. Abbreviations: CVRM = cardiovascular risk management, GDM = gestational diabetes mellitus, HDP = hypertensive disorders of pregnancy, CVD = cardiovascular disease, GP = general practitioner.

and monitoring after reproductive disorders, the central CVRM GP guideline does not. According to GPs, this leads to increased practice variation. Finally, CVRM is not structurally covered after HDP/GDM. HCPs indicate this hampers implementation and innovation. HCPs further noted a lack of regional protocols and interdisciplinary communication at hospital discharge.

To improve current CVRM, HCPs emphasized the importance of interdisciplinary communication across settings to ensure continued CVRM. According to them, this may benefit groups already at higher risk due to lower SES and health literacy. They suggested creating regional protocols with standardized phrases in referral letters to improve care continuity and communication. Midwives suggested a routine week 6-

appointmentfor those that did not give birth in primary care. Moreover, GPs noted that better identifiability of patients in records is needed to allow active outreach and routine monitoring.

3.2. Preferences for CVD risk education, lifestyle interventions and routine GP monitoring

3.2.1. CVD risk education

Both women and HCPs acknowledged that being informed about CVD risk is a first, essential step for women to engage in CVRM. According to women, being informed allows them to try and prevent CVD in the future and prepare their health for future pregnancies (Table 2,

quote 5). While women from all backgrounds expressed the desire to receive CVD risk education from their HCPs, a few preferred not to be informed during the pregnancy because of potential added stress. In general, repeated information provision was desired during and beyond pregnancy, for example at diagnosis, at the week 6-appointment by the obstetrician or community midwife, and after 3–6 months by the GP. Overall, the week 6-appointment was mentioned by women in all groups as the most appropriate moment for CVD risk education, even by those who missed the appointment. However, even when a week 6-appointment was planned, women and HCPs indicated it is currently underutilized for the purpose of CVRM because it centers around postpartum recovery. In general, women prefer to receive information in-person from a trusted HCP (Table 2, quote 6), so they can ask questions. Furthermore, women mentioned that information leaflets are likely to go unread although there was some heterogeneity between women depending on ethnic background. Only women that are highly educated mentioned a preference for websites. HCPs added that information on future health risks should at least be given before the patient is released from obstetric care while allowing women time to adapt to their new circumstances (Table 2, quote 7). HCPs confirmed that leaflets should not be the only source of CVD risk education.

3.2.2. Lifestyle interventions

Around half of the women indicated that they were open to personalized lifestyle interventions. The preferred timing of lifestyle interventions according to women, including specific support for weight loss, was around 6 months, and initiated by the GP (or dietician, physical therapist, exercise program). Women also mentioned that putting recommendations into practice is challenging and they need help doing so (Table 2, quote 8). Women who had a negative attitude towards lifestyle interventions were often beyond four years postpartum. We were unable to discern if this was because of lowering of CVD risk perception or lost motivation. Those working in healthcare perceived themselves to be self-sufficient in making the required lifestyle changes.

HCPs from hospital settings recognize that they have limited time (up to six weeks postpartum and in short consultations) to establish behavioral changes and therefore focus on improvements during pregnancy. They feel the designated place for continued lifestyle interventions is in the GP practices. Some HCPs proposed the use of lifestyle apps, but only highly educated women actively mentioned using them. Additionally, HCPs emphasized that effectively changing lifestyle behavior requires motivational counselling skills and tailored interventions to suit patient's personal situation and preferences. All HCPs mentioned ethnic background needs to be taken into account.

Lastly, according to HCPs solutions should not just be sought in healthcare. They advocate for more general measures, such as greater societal awareness of hypertension and weight management (Table 2, quote 9).

3.2.3. Routine GP monitoring

Most women were positive about routine blood pressure and glucose value checks, perceiving such monitoring as important for their health, an opportunity to discover CVD risk factors sooner and be a healthy mother. However, a few women mentioned that out-of-pocket costs (e.g. for laboratory tests), as a barrier to routine monitoring.

In all women, frequently emerging theme was the desire to be actively invited and reminded by the GP. Women indicated this would be a solution to mothers prioritizing other things over their own health and therefore often do not initiate these appointments (Table 2, quote 10). This finding seemed not so much related to the number of children women had, but to being a mother and performing the role of caretaker. Sometimes, women had received opportunistic monitoring during visits for other health problems (i.e., iron deficiency, contraception) and found that this was an efficient way of receiving monitoring. Additionally, some women said they would like a stronger focus on lifestyle interventions (which would perceptively normalize the future risk of CVD)

than on routine GP monitoring. These women were all highly educated or healthcare providers.

To facilitate early detection and outreach, and to lower financial barriers for women, some GPs and GP practice nurses suggested to include women after HDP/GDM in the Dutch integrated care program for CVRM for routine monitoring. Other GPs were hesitant to add groups to the Dutch integrated CVRM program due to ongoing debate among GPs about the growing focus on prevention, potentially overburdening the primary care system. Several HCPs called for strengthening of regional protocols and collaborations. As an example of collaboration, we found GPs that organized their clinics to be highly integrated with a community midwife clinic. This initially served a practical purpose: to prevent mental health problems and unintended pregnancies. Due to the close collaboration, the involved HCPs perceived they could keep better track of women who had HDP/GDM and provide CVRM.

3.3. Focus points for postpartum-CVRM in a multi-ethnic population

A general pattern was observed of women with an ethnic minority background, lower SES and lower disease severity being less informed and more often missing the week 6-appointment, routine GP monitoring, and the potential subsequent lifestyle interventions. Women with an ethnic minority background often reflected on the downstream effects of missing important health information (sometimes not through their own direct experience but observed in members of their family or community). This was not mentioned by the majority group. Unclear information and recommendations may lead to symptoms not being recognized and delayed care. This may be further compounded when people do not understand the healthcare system (Table 2, quote 11), and by cultural perceptions of pregnancy which may influence lifestyle behavior that can persist in the postpartum period. Although women in general preferred in-person information transfer to written information, written material in several languages was desired to reach people with a language barrier. Women suggested that to adequately reach people with ethnic minority backgrounds, information transfer should take place in people's own environments, like community gatherings, outside of the medical system. Moreover, women with ethnic minority backgrounds more often spontaneously mentioned the relevance of having family members with diabetes or CVD who actively supported them with advice (e.g., to reduce salt intake, home blood pressure monitoring and routine monitoring). At the same time, family members had a negative influence when the family members felt it was a burden to adjust dietary patterns to their needs, dismissed their health concerns or implied they were to blame for the pregnancy complications. Lastly, it was noted that several women with an ethnic minority background experienced being stereotyped as unhealthy, not being provided with adequate care, or being looked down upon by their HCPs because of their ethnic background or language difficulties.

The need for understandable risk communication to people with diverse backgrounds was also a theme that arose frequently among HCPs. HCPs explained that when it is challenging to deliver complex health information, they often limit their communication to more acute health problems. HCPs suggested using translators (translation service or a family member) and providing translated written material to bridge language barriers. However, translated written material on this specific topic is often not available. HCPs also indicated that health- and lifestyle recommendations are often not applicable in ethnically diverse populations that may have their own dietary and physical activity preferences. To motivate women and make recommendations easier to adhere to, personal tailoring and targeting to specific needs of certain populations is perceived as essential and was mentioned by all HCPs (Table 2, quote 12). Lastly, HCPs highlight that what makes people vulnerable may be less tied to ethnicity or having an ethnic minority background, but more often to language barriers, lower SES, health literacy, higher parity, or other health concerns like having mental health problems (Table 2, quote 13).

4. Discussion

4.1. Principal findings

Regardless of ethnic background, our study showed a high drop out from the CVRM care pathway after pregnancies complicated by HDP/GDM, particularly at the transfer from obstetric to primary care and before long-term routine monitoring. Women were frequently unaware of the relationship between HDP/GDM and their subsequent increased risk of CVD and had not been followed up. This pattern occurred more often in groups with an ethnic minority background and/or lower SES. Women want to receive well-timed education and lifestyle advice, in-person from their HCPs, that is targeted to their specific (cultural) circumstances, as well as active outreach for monitoring. HCPs signal the need to resolve system barriers and implementation of more structured interdisciplinary communication across settings supported by regional protocols and collaborative care organization.

4.2. Results in the context of what is known

Our general findings align with previous international literature describing a lack of awareness of CVD risk after HDP/GDM among women and HCPs, with women from all backgrounds expressing a desire for CVD risk education, personalized lifestyle interventions, and extra check-ups [11,12,14,15]. Our investigation adds to previous work in several ways: we showed that the short-term aftercare, specifically the week 6-appointment, is currently underutilized for CVD risk education and is a major point of dropout. By patient journey mapping and contrasting the experienced and perceived care trajectories across primary, secondary, and tertiary settings, we identified this critical juncture in CVRM.

We also identified specific needs and barriers for women from ethnic minorities, due to our larger and more socioeconomically diverse multi-ethnic sample, e.g., the concerns around out-of-pocket cost. Additionally, we show the need for culturally tailored information and lifestyle interventions, which Joo et al. (2021) found can reduce health disparities and improve care quality for ethnic minorities [24]. However, care must be taken to avoid stereotyping or discrimination, which women in our study reported experiencing. Zemouri et al. (2024) recently showed that people with ethnic minority backgrounds experience discrimination in Dutch healthcare [25,26]. Our study further offers broader insights into HCP perspectives, revealing that while most HCPs recognize the potential benefit, there is much variation in GP support for routine CVRM after HDP/GDM, with barriers listed varying from reimbursements to a perceived lack of evidence on effectiveness.

4.3. Clinical implications

We identified several leverage points to improve the provision and uptake of postpartum-CVRM being; active outreach, the timely in-person delivery of CVD risk education, with translated written materials in several languages, and personally tailored (culturally sensitive) lifestyle interventions. Specifically, within Dutch context, a routine week 6-appointment has been suggested by midwives in our study, which would ensure women (who transferred across settings) get information about their future health and monitoring. However, our findings suggest that until these measures are uniformly included in all guidelines and regional protocols, implementation may remain limited.

4.4. Research implications

Previous studies have proposed including women after HDP/GDM in the GP CVRM guideline and the Dutch (insurance covered) integrated care program [12,13]. Our study supports that this may improve uptake and the continuation of CVRM and reduce barriers that are sources of disparities. This may require adopting a 'life course approach' to CVD

risk in women [26] and gaining more broad support among GPs. To increase support, we recommend further research efforts to focus on generating evidence on the effectiveness of the prevention of CVD risk factors after HDP/GDM.

4.5. Strengths and limitations

There are several study limitations to consider that have impacted our findings. First, we did not achieve the desired quota of 6–8 participants in all ethnic subgroups, although the composition of our patient and HCP samples reflects our effort to recruit a diverse group of women and HCPs. This means we were not able to compare subgroups by ethnicity.

Second, we are cautious to generalize our findings to populations outside of our study sample and context, which is a general concern in qualitative research. Our findings primarily reflect the experiences of individuals from diverse ethnic backgrounds who were born or raised in the Netherlands, rather than those of first-generation migrants or groups with limited language proficiency. The ethnic background of the interviewers (e.g., Dutch, Moroccan) may have facilitated recruitment in the groups with Dutch and Moroccan backgrounds within the project duration, while lower number were achieved for other groups. Interviewers proficient in additional languages than Dutch and Darija (Moroccan Arabic), could have improved our ability to reach these women. Moreover, our findings may not reflect experiences across all SES groups; our final sample had a higher proportion of women with higher education in the Dutch group and more unemployed women in the ethnic minority group. In addition, women from the lowest SES groups were underrepresented in all groups. We have tried to mitigate this by asking participants (both patients and HCPs) about their views on how women with diverse ethnic and SES backgrounds, also those not easily reached for research, could best be supported. Although this gave us valuable insights, it cannot replace accounts of direct experiences. Thus, despite reaching conceptual saturation in our analysis, adding more women with specific characteristics might have yielded additional insights.

Further, the generalizability of our findings might have been improved if we had been able to include a larger group of HCPs. By design, the HCPs subgroups were too small to reach saturation.

Finally, because we focused on a single context, our findings may reflect experiences, barriers, and improvements specific to the Dutch healthcare context. In the Dutch healthcare system obstetrical care is differently organized than in other countries. Moreover, Dutch primary care has a leading role in long-term CVRM in high CVD risk populations. This may limit the applicability of our findings to other settings or underestimate the disparities in countries with different organization of birth care and subsequent CVRM, such as the United States.

5. Conclusions

In conclusion, our study highlights that, at present, there is a high rate of drop out for CVRM after HDP/GDM in women with diverse ethnic and socioeconomic backgrounds in the Netherlands. Leverage points to mitigate dropout and improve postpartum CVRM were identified, including active outreach by healthcare providers to ensure continuation of care, better utilization of existing follow-up moments for CVD risk education and offering culturally sensitive support. Additionally, removing the system barriers, more evidence-based guidelines and implementation of regional multidisciplinary protocols could reduce barriers for healthcare providers to provide continued CVRM and reduce health disparities in increasingly diverse settings.

Author contributions

RB, WG, SG, and IvV conceived and designed the HELIUS-Perined study project. AR drafted this specific study and all study components,

including obtaining medical ethical approval, developing topic lists and recruitment materials. AR served as the executive researcher, collecting, and analyzing data, and writing the first draft of the manuscript. SB contributed to data collection and analysis specifically for the patient group with a Moroccan background. SMC provided methodological expertise and supported the integrated patient journey mapping. IvV supervised AR throughout the process. All authors reviewed and provided feedback on the final manuscript and agree with its content.

Funding

This study was part of the HELIUS-Perined research project and was funded by ZonMw (grant numbers 849,200,008 and 05430052110005). The funding source had no role in study design; in data collection, analysis, and interpretation; in the report's writing; and the decision to submit the article for publication.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

We would like to thank all the participants for contributing their time and efforts, both women and health care providers, without whom this study would not have been possible. Next, we are grateful for the support we received from the Dutch HELLP foundation and the regional obstetric care consortia of the North and North-West Netherlands and their communication for recruitment and dissemination. We also thank A. Moro, M. Wagemakers and B. Leijendekker (students of the academy for midwifery in Amsterdam) for transcribing part of the interviews.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.preghy.2025.101203>.

References

- [1] L. Benschop, J.J. Duvekot, J.E.R.v. Lennep, Future risk of cardiovascular disease risk factors and events in women after a hypertensive disorder of pregnancy, *Heart* 105 (16) (2019) 1273–1278.
- [2] H.L. Brown, G.N. Smith, Pregnancy complications, cardiovascular risk factors, and future heart disease, *Obstetrics and Gynecology Clinics*. 47 (3) (2020) 487–495.
- [3] C.K. Kramer, S. Campbell, R. Retnakaran, Gestational diabetes and the risk of cardiovascular disease in women: a systematic review and meta-analysis, *Diabetologia* 62 (6) (2019) 905–914.
- [4] G.N. Smith, J.M. Louis, G.R. Saade, Pregnancy and the postpartum period as an opportunity for cardiovascular risk identification and management, *Obstet. Gynecol.* 134 (4) (2019) 851–862.
- [5] J. Lewey, T.M. Beckie, H.L. Brown, S.D. Brown, V.D. Garovic, S.S. Khan, E. C. Miller, G. Sharma, L.S. Mehta, American Heart Association cardiovascular disease and stroke in women and underrepresented populations committee of the council on clinical cardiology; council on cardiopulmonary, critical care, perioperative and resuscitation; and council on cardiovascular and stroke nursing. opportunities in the postpartum period to reduce cardiovascular disease risk after adverse pregnancy outcomes: a scientific statement from the American heart association, *Circulation* 149 (7) (2024) e330–e346.
- [6] CVRM na een reproductieve aandoening - Federatie Medisch Specialisten 2023 [Available from: https://richtlijnendatabase.nl/richtlijn/cvrm_na_een_reproductieve_aandoening/cvrm_na_reproductieve_aandoening_-_startpagina.html. Accessed August 30, 2024.
- [7] D.T. Gamble, B. Brikinns, P.K. Myint, S. Bhattacharya, Hypertensive disorders of pregnancy and subsequent cardiovascular disease: current national and international guidelines and the need for future research, *Front. Cardiovasc. Med.* 17 (6) (2019) 55.
- [8] E.W. Seely, A.C. Celi, J. Chausmer, C. Graves, S. Kilpatrick, J.M. Nicklas, et al., Cardiovascular health after preeclampsia: patient and provider perspective, *J. Womens Health (Larchmt)* 30 (3) (2021) 305–313.
- [9] D. Bick, S.A. Silverio, A. Bye, Y.-S. Chang, Postnatal care following hypertensive disorders of pregnancy: a qualitative study of views and experiences of primary and secondary care clinicians, *BMJ Open* 10 (1) (2020) e034382.
- [10] F. van Kesteren, S. Visser, W. Hermes, P.W. Teunissen, A. Franx, M.G. van Pampus, et al., Counselling and management of cardiovascular risk factors after preeclampsia, *Hypertens. Pregnancy* 35 (1) (2016) 55–61.
- [11] H.L. Sandsaeter, J. Horn, J.W. Rich-Edwards, H.S. Haugdahl, Preeclampsia, gestational diabetes and later risk of cardiovascular disease: women's experiences and motivation for lifestyle changes explored in focus group interviews, *BMC Pregnancy Childbirth* 19 (1) (2019) 448.
- [12] T.E. Dijkhuis, F. Bloem, L.A.J. Kusters, S.M. Roos, S.J. Gordijn, F. Holvast, J. R. Prins, Investigating the current knowledge and needs concerning a follow-up for long-term cardiovascular risks in Dutch women with a preeclampsia history: a qualitative study, *BMC Pregnancy Childbirth* 20 (1) (2020) 486.
- [13] S. van Smoorenburg, J.M. Kist, R.C. Vos, H.M.M. Vos, Experiences with an integrated screening programme targeted at women who had a hypertensive disorder or diabetes in pregnancy in the Netherlands: a qualitative study, *BMJ Open* 13 (8) (2023) e066198.
- [14] J.A. E. van Oortmerssen, B. Hummel, E. Aribas, A. Wijtzes, S. Polinder, I. Spronk, J. J. Duvekot, J.E. Roeters van Lennep, On behalf of the IMPRESS Consortium, Tailoring lifestyle guidance for women after pregnancies complicated by severe pre-eclampsia: a mixed-methods study into women and healthcare provider perspectives, *European Journal of Preventive Cardiology*, 2024; zwae232.
- [15] G. Skurnik, A.T. Roche, J.J. Stuart, J. Rich-Edwards, E. Tsigas, S.E. Levkoff, E. W. Seely, Improving the postpartum care of women with a recent history of preeclampsia: a focus group study, *Hypertens. Pregnancy* 35 (3) (2016) 371–381.
- [16] E. Chan, C.M. Nowik, J. Pudwell, G.N. Smith, Standardized postpartum follow-up for women with pregnancy complications: barriers to access and perceptions of maternal cardiovascular risk, *J. Obstet. Gynaecol. Can.* 43 (6) (2021) 746–755.
- [17] R.J. Burger, S.J. Gordijn, R. Boliijn, A. Reilingh, E.P. Moll Van Charante, B.H. Van Den Born, et al., Cardiovascular risk profile after a complicated pregnancy across ethnic groups: the HELIUS study, *Eur. J. Prev. Cardiol.* (2022).
- [18] W. Perini, A.E. Kunst, M.B. Snijder, R.J.G. Peters, I.G.M. van Valkengoed, Ethnic differences in metabolic cardiovascular risk among normal weight individuals: implications for cardiovascular risk screening. the HELIUS study, *Nutr. Metab. Cardiovasc. Dis.* 29 (1) (2019) 15–22.
- [19] C. Agyemang, B.J. van den Born, Cardiovascular health and disease in migrant populations: a call to action, *Nat. Rev. Cardiol.* 19 (1) (2022) 1–2.
- [20] S. McCarthy, P. O'Raghallaigh, S. Woodworth, Y.L. Lim, L.C. Kenny, F. Adam, An integrated patient journey mapping tool for embedding quality in healthcare service reform, *J. Decis. Syst.* 25 (sup1) (2016) 354–368.
- [21] S. McCarthy, P. O'Raghallaigh, S. Woodworth, Y.Y. Lim, L.C. Kenny, F. Adam, Embedding the pillars of quality in health information technology solutions using "integrated patient journey mapping" (IPJM): case study, *JMIR Hum. Factors* 7 (3) (2020) e17416.
- [22] V. Braun, V. Clarke, Using thematic analysis in psychology, *Qual. Res. Psychol.* 3 (2) (2006) 77–101.
- [23] B.C. O'Brien, I.B. Harris, T.J. Beckman, D.A. Reed, D.A. Cook, Standards for reporting qualitative research: a synthesis of recommendations, *Acad. Med.* 89 (9) (2014) 1245–1251.
- [24] J.Y. Joo, M.F. Liu, Culturally tailored interventions for ethnic minorities: a scoping review, *Nurs. Open* 8 (5) (2021) 2078–2090.
- [25] C. Zemouri, A. Nait Kassi, W. Arrazola de Oñate, et al., Exploring discrimination and racism in healthcare: a qualitative phenomenology study of Dutch persons with migration backgrounds, *BMJ Open* 2024;14:e082481.
- [26] G. Mishra, R. Hardy, D. Kuh (eds), *A Life Course Approach to Women's Health*, 2nd edn (Oxford, 2023; online edn, Oxford Academic, 20 July 2023).