

Inadequate Health Literacy in Patients with Arterial Vascular Disease

Ruben M. Strijbos^{*}, Jan-Willem Hinnen, Ronald F.F. van den Haak, Bart A.N. Verhoeven, Olivier H.J. Koning

Jeroen Bosch Hospital Netherlands, 's-Hertogenbosch, The Netherlands

WHAT THIS PAPER ADDS

Knowledge of the health literacy of the patient population is important in the context of supplying medical information, shared decision making, and improving care and health outcomes. Patients with arterial vascular disease are a large subset of the population visiting the vascular surgeon with specific characteristics that are associated with inadequate health literacy. This is the first large scale cross sectional study providing data about the prevalence of inadequate health literacy in patients with arterial vascular disease.

Objective: The aim was to identify the prevalence of inadequate health literacy in patients with arterial vascular disease. This was a cross sectional study.

Methods: Patients with arterial vascular disease visiting the outpatient clinic between January 5, 2015 and December 28, 2016, were randomly included and screened for inadequate health literacy with the Newest Vital Sign—Dutch (NVS-D), a validated health literacy assessment measure. A score of <4 out of six identified individuals with inadequate health literacy. Age, gender, highest education level, and reason for consultation were also registered. Data analysis was performed using Student's *t*-test or the Mann–Whitney *U* test and chi-square test. Logistic regression with backward elimination was applied to identify independent predictors.

Results: A total of 202 patients were included. The mean NVS-D score was 1.91 (SD $\pm$ 1.948, median 1). The prevalence of inadequate health literacy was 76.7%. A significantly higher prevalence of inadequate health literacy was found in patients ≥ 65 years ($p < .001$) and patients with a lower education level ($p < .001$). No significant difference was found between female/male patients ($p = .056$), nor between participants with peripheral arterial occlusive disease and abdominal aortic aneurysm ($p = .116$). Age (OR 1.060; 95% CI 1.017–1.104; $p = .005$) and education level (OR 0.164; 95% CI 0.078–0.346; $p < .001$) were identified as independent predictors of inadequate health literacy.

Conclusion: This study shows a prevalence of inadequate health literacy of 76.7% in patients with arterial vascular disease, with a significantly higher prevalence in patients ≥ 65 years and patients with a lower education level. The high prevalence of inadequate health literacy should be considered when information is provided, and suggests the need to further investigate the best methods to convey medical information to this group of vulnerable patients.

© 2018 European Society for Vascular Surgery. Published by Elsevier B.V. All rights reserved.

Article history: Received 24 May 2017, Accepted 22 April 2018, Available online 8 June 2018

Keywords: Health literacy, Peripheral arterial disease, Abdominal aortic aneurysm, Patient education

INTRODUCTION

Despite modern education, low literacy skills are still a common problem in The Netherlands and other parts of Europe and the United States.^{1–3} A recently published international survey, the PIAAC (Program for the International Assessment of Adult Competencies), shows that in The Netherlands 1.3 million adults between 18 and 65 years (approximately 30%) have low literacy skills.¹ The ability to write and read is an important condition for adequate health literacy.⁴ Health literacy is defined as the degree to

which individuals have the capacity to obtain, process, and understand basic health information and services needed to make appropriate health decisions. Every medical situation requires several generic or specific skills, so the term health literacy consists of different capabilities, for example reading and numeracy skills.⁵

Knowledge about the health literacy of a patient population is important in the context of providing medical information in a comprehensible way, shared decision making, and increasing patient satisfaction. Moreover, several studies have already shown that inadequate health literacy is consistently associated with poorer health outcomes and poorer use of healthcare services.^{6,7} Also, different studies show evidence of the association between low health literacy and adverse clinical events and increased healthcare use in specific patient groups, such as patients

^{*} Corresponding author. Jeroen Bosch Hospital Netherlands, Postbus 90153, 5200 ME 's-Hertogenbosch, The Netherlands.

E-mail address: r.strijbos@jbz.nl (Ruben M. Strijbos).

1078-5884/© 2018 European Society for Vascular Surgery. Published by Elsevier B.V. All rights reserved.

<https://doi.org/10.1016/j.ejvs.2018.04.015>

with chronic kidney disease⁸ and coronary artery disease.⁹ Furthermore, a current meta-analysis revealed the importance of health literacy and the efficacy of health literacy interventions in adherence to medical treatment in chronic and acute illness.¹⁰ Nevertheless to date, relatively little is known about health literacy among the general population in Europe.

Patients with arterial vascular disease are an important and specific subset of the population visiting the department of vascular surgery. Arterial vascular disease is linked to diabetes mellitus,¹¹ smoking,¹² ageing,¹³ lower socio-economic status and lower education level.¹⁴ Previous studies have shown that inadequate health literacy is associated with these factors as well.^{15–18} The relationship between health literacy and ethnic/racial disparities on health outcomes remains rather limited and mixed.¹⁹ The hypothesis is that inadequate health literacy may be a common problem in the group of patients with arterial vascular disease. There is no literature describing the extent of the problem of inadequate health literacy in patients with arterial vascular disease. Understanding this problem and its magnitude could influence the way information is provided. The aim of this study was to identify the prevalence of inadequate health literacy in patients with arterial vascular disease.

METHODS

In this cross sectional study, patients with established peripheral arterial occlusive disease (PAOD) or an abdominal aortic aneurysm (AAA) were included between January 5, 2015 and December 28, 2016. The first cohort included the period between January 5, 2015 and January 16, 2015. In addition, a second cohort with unique patients was included spread over a period between April 21, 2016 and December 30, 2016, in order to create a larger and more representative sample. There was a total of 21 inclusion days in this aforementioned period of time. The researchers scheduled their inclusion dates without of the knowledge of the staff at the outpatient clinic. Moreover, patients were randomly selected by the researchers from a list of names at the outpatient clinic (without access to their medical charts or any other patient information). These precautions were taken in order to create a random sample. There was no formal randomisation.

All patients, both initial consults and follow up contacts, were visiting the outpatient clinic of the department of vascular surgery in a large Dutch teaching hospital. Exclusion criteria were a recent stroke (within 1 year), congenital or traumatic brain damage, and blindness or deafness. People who needed an interpreter to communicate were also excluded. After the appointment with their surgeon, patients were asked to participate in the study.

The diagnosis of arterial vascular disease in patients was established in accordance with the applicable (inter)national guidelines. Peripheral arterial occlusive disease was diagnosed in patients with an abnormal ankle brachial index in rest (<0.9) or a drop of the ankle brachial index of >0.15 during exercise. An abdominal aortic aneurysm was confirmed when there was focal dilatation of the blood

vessel that was 50% larger than the normal aortic diameter on computed tomography angiography (CTA) or ultrasound.

The validated Dutch version of the widely used US and UK Newest Vital Sign (NVS),²⁰ the Newest Vital Sign—Dutch (NVS-D), was used as screening instrument. This short questionnaire consisted of six brief questions, which the patient had to answer orally. A score of <4 out of 6 identified individuals with inadequate health literacy, because this cut off point distinguished best between individuals with adequate versus inadequate health literacy.²¹ The original (English) version of the Newest Vital Sign is shown in Fig. 1. Age, gender, highest education level, and reason for consultation were also ascertained as background characteristics. Finally, the medical charts of the patients were reviewed for smoking, diabetes, ischaemic heart disease, and hypertension as characteristics of the population.

The analysis of health literacy and NVS-D scores, in the context of the patient characteristics, was performed with the Student *t*-test if there was a normal distribution or the Mann–Whitney *U* test if there was no normal distribution. To determine if there was a normal distribution, the Kolmogorov–Smirnov test was used. If the outcome was a proportion, the chi-square test was performed or the Fisher exact test if appropriate. The level of significance applied was $p = .05$. In order to identify independent predictors for inadequate health literacy, a logistic regression with backward elimination was applied. Backward selection started with all of the variables (age, gender, education level, and reason for consultation; i.e., abdominal aortic aneurysm or peripheral arterial occlusive disease) in the model, and the variables were removed one by one as they were found to be insignificant in predicting the outcome. A backward selection (starting with all covariates) was performed with a significance level of $\alpha = 0.10$ for staying in the model; all factors significant at $p \leq .10$ were retained in the final model. The goodness of fit (Hosmer and Lemeshow test), adjusted odds ratios, and 95% confidence interval were calculated from the final logistic regression model. All analyses were performed using Statistical Package for Social Sciences (IBM SPSS Statistics for Windows, Armonk, NY; IBM Corp), version 23.0.

Ethics

The regional Medical Ethics Committee stated that this study did not require formal approval of the Netherlands Medical Research Involving Human Subjects Act. All potential respondents were informed about the aim of the study by a written form and orally, with implementation of interventions for patients with limited health literacy skills (see also the Discussion). The participation was voluntary. The potential respondents could opt out of the study at any time. By participating, respondents consented to the research procedure.

RESULTS

Patients

A total of 202 patients with arterial vascular disease were included. There were seven patients who declined

Table 1. Characteristics of the study participants.

Background characteristics	n (202)	%
Age, years		
<55	11	5.4
55–64	47	23.3
65–74	77	38.1
75–84	51	25.2
≥85	16	7.9
Gender		
Male	122	60.4
Female	80	39.6
Highest education level		
None	3	1.5
Elementary school	56	27.7
Secondary school	63	31.2
Intermediate vocational education	35	17.3
Higher vocational education	30	14.9
University	7	3.5
Unknown	8	4.0
Arterial disease		
Peripheral arterial occlusive disease	147	72.8
Abdominal aortic aneurysm	55	27.2
Smoking^a		
Yes	97	48.0
No	105	52.0
Diabetes^b		
Yes	59	29.2
No	143	70.8
Ischaemic heart disease^c		
Yes	61	30.2
No	141	69.8
Hypertension^d		
Yes	133	65.8
No	69	34.2

^a The criterion for smoking was the use of tobacco products at the time of consultation.

^b Diabetes mellitus was diagnosed conforming to (inter)national applicable standard: fasting plasma glucose ≥ 7.0 mmol/L (126 mg/dL) or 2 h plasma glucose ≥ 11.1 mmol/L (200 mg/dL).

^c Ischaemic heart disease was registered in case of angina pectoris or (an episode of) myocardial ischaemia.

^d Hypertension was defined as a systolic blood pressure of 140 mmHg or more or a diastolic blood pressure of 90 mmHg or more.

patient satisfaction and more favourable health outcomes if clearer medical information is provided.^{23,24}

These findings imply the need to look at the medical information supply for patients with arterial vascular disease in a critical way. Recently, a large study was performed with the aim of determining the prevalence of limited health literacy among pharmacy visitors in The Netherlands. Almost 1000 adult participants were included and the NVS-D was also used to identify inadequate health literacy. This study found a prevalence of limited health literacy skills of 52%.²⁵ In comparison with the prevalence of 76.7% in this population of patients with arterial disease, this only emphasises the need to further investigate the best methods to convey (medical) information.

In the literature, some research has already been performed to discover interventions for patients with inadequate health literacy. A recent large systematic review of Sheridan et al.²⁶ reviewed 38 studies and focused on the effectiveness of interventions for those with low health literacy. Some examples are presenting essential information concisely or first on the paper, presenting information so that a higher number is better (i.e. a higher number indicating better quality)²⁷ and the addition of a verbal narrative in instructional videos.²⁸ Intensive mixed strategy interventions focusing on self management led, in some of the included studies, to better use of health care services.^{29–34} Still, it can be difficult to integrate these self management interventions in daily practice. Next to these promising results, not all interventions with intuitive appeal are necessarily successful. For instance, four included randomised controlled trials and one other prospective study evaluated the effects of substituting diverse media for print and found contradictory results. For example, Meade et al.³⁵ found no differences between written and video-taped information. In contrast, Murphy et al.³⁶ described that providing information by video alone may have limited benefits, while Bryant et al.³⁸ noticed a significant difference in favour of multimedia versions of patient information.^{35–39}


**Figure 2.** Scores on the Newest Vital Sign — Dutch.

Table 2. Health literacy in the context of patient characteristics.

	Inadequate Health Literacy		Adequate Health Literacy		<i>p</i>
	n	%	n	%	
Total patients	155	76.7	47	23.3	
Gender					
Male	88	72.1	34	27.9	.056
Female	67	83.8	13	16.3	
Age, mean, years ($\pm$ SD)	71.2 ($\pm$ 9.404)		65.6 ($\pm$ 9.046)		< .001
Range	45–95		41–89		
< 65 years	35	60.3	23	39.7	< .001
$\geq$ 65 years	120	83.3	24	16.7	
Education					
Low	108	88.5	14	11.5	< .001
High	40	55.6	32	44.4	
Arterial disease					
Peripheral arterial occlusive disease	117	79.6	30	20.4	.116
Abdominal aortic aneurysm	38	69.1	17	30.9	

The education level was unknown in eight patients.

Bold *p* values are significant differences ($p < .05$).

An article by Hironaka et al.⁴⁰ explored strategies to improve patient–provider communication for clinicians who care for people with limited health literacy. In the context of verbal communication, some suggestions are communication using plain language, avoiding jargon, limiting the items discussed during a consultation and the use of multiple forms of communication (such as a combination of pictures, written, and oral information). Another recent study showed that spoken animation is the best way to communicate complex information to patients with low health literacy skills.⁴¹ Although some advances have been made in the field of discovering methods for effective information supply for patients with low health literacy in the last decade, further well-designed research should be performed in order to discover the best method of information supply and to support evidence for the current strategies. Preferably, research should also focus on specific patient populations, like patients with arterial vascular disease because of the high prevalence of the problem in this subset of patients.

The current study has several strengths, besides the previously mentioned prospective study design. Firstly, over

200 patients were included. The participation was minimally invasive and took only about 5 min for patients, so there were only few patients who declined participation. Secondly, only three well instructed researchers screened for health literacy with the NVS-D in all patients and there were no missing data (except the education level of 8 patients). Finally, the analysis of the differences in inadequate health literacy in the context of age and education level showed the same association as stated in previous studies in the general population.^{17,18}

A limitation of this study is the screening instrument used, the NVS-D. The NVS-D could not differentiate the degree of inadequate health literacy. It indicates only if a patient has adequate or inadequate health literacy. Also, it measures the objective health literacy and does not give any information regarding the subjective health literacy: knowledge about the self reported confidence or possible difficulties with understanding information patients encounter in the health care setting. The available Dutch test for subjective health literacy is a version of the Set of Brief Screening Questions (SBSQ).⁴² However, the Dutch version of the SBSQ was not validated at the time of this research.⁴³ In addition, the NVS-D itself does have some limitations in identifying health literacy. By using the cut off point <4 to identify inadequate health literacy, the NVS-D would correctly identify 80% of those with adequate health literacy and 56% of those with inadequate health literacy in Dutch patients. Nevertheless, only very few Dutch screening instruments for health literacy are available and the mentioned cut off scores were determined as the most optimal by the receiver operating characteristics (ROC) in the validation study.²¹ Furthermore, a short Dutch screening tool for illiteracy is not available. That information could have given more insight into the interpretation of inadequate health literacy in this patient population.

Another possible confounding factor was the timing of the test. The participants completed the NVS-D after their appointment with the vascular surgeon. In this way, the study did not interfere with the schedule at the outpatient

Table 3. Mean NVS-D scores in relation to patient subgroups.

	Mean NVS-D score ($\pm$ SD)	<i>p</i>
Gender		
Male	2.16 (2.074)	.061
Female	1.53 (1.676)	
Age		
<65 years	2.64 (2.108)	.01
$\geq$ 65 years	1.62 (1.805)	
Education		
Low	1.29 (1.582)	< .001
High	3.07 (1.974)	
Arterial disease		
Peripheral arterial occlusive disease	1.78 (1.842)	.218
Abdominal aortic aneurysm	2.27 (2.181)	

Bold *p* values are significant differences ($p < .05$).

clinic and patients did not have to come to the hospital at another time to complete the test. The timing of the test after the appointment may have had a negative effect on the concentration and motivation of the patients, influencing the scores on the NVS-D negatively. On the other hand, this is the moment in the consultation where medical information is usually given to the patient and therefore represents the regular mindset of patients within the information process.

Finally, it is important to consider that low literacy skills and illiteracy are still taboo subjects.⁴⁴ A lot of people with inadequate literacy skills are embarrassed about this problem and will try to avoid difficult situations in which they can be confronted with it. In this way, the social taboo undoubtedly has an influence on research into the field of health literacy.

CONCLUSION

This study shows a prevalence of inadequate health literacy of 76.7% in patients with arterial vascular disease, with a significantly higher prevalence in the subgroups of patients ≥ 65 years and patients with a lower education level. The high prevalence of inadequate health literacy should be considered when information is provided and suggests the need to further investigate the best methods to convey medical information to this group of vulnerable patients.

ACKNOWLEDGEMENTS

The authors would like to thank W. Koning and T. van Bouwdijk for their contribution in screening patients with the NVS-D and Dr. M. Hilbink for her help in the statistical analyses.

CONFLICT OF INTEREST

None.

FUNDING

None.

REFERENCES

- Buisman M, Allen J, Fouarge D, Houtkoop W, Van der Velden R. *PIAAC: kernvaardigheden voor werken en leren, resultaten nederlandse survey 2012*. 's-Hertogenbosch: ECBO; 2013.
- Kutner M, Greenberg E, Jin Y, Boyle B, Hsu Y, Dunleavy E. *Literacy in everyday life: results from the 2003 national assessment of adult literacy*. Washington, DC: National Center for Education Statistics; 2007.
- OECD. *OECD: skills outlook 2013: first results from the survey of adult skills*. Paris: OECD Publishing; 2013.
- Andrus MR, Roth MT. Health literacy: a review. *Pharmacotherapy* 2002;**22**:282–302.
- Nielsen-Bohlman L, Panzer AM, Kindig DA. *Health literacy: a prescription to end confusion*. Washington, DC: The National Academy Press; 2004.
- Berkman ND, Sheridan SL, Donahue KE, Halpern DJ, Crotty K. Low health literacy and health outcomes: an updated systematic review. *Ann Intern Med* 2011;**155**:97–107.
- Dewalt DA, Berkman ND, Sheridan S, Lohr KN, Pignone MP. Literacy and health outcomes: a systematic review of the literature. *J Gen Intern Med* 2004;**19**:1228–39.
- Taylor DM, Fraser S, Dudley C, Oniscu GC, Tomson C, Ramanan R, et al. Health literacy and patient outcomes in chronic kidney disease: a systematic review. *Nephrol Dial Transpl* 2017. <https://doi.org/10.1093/ndt/gfx293> [Epub ahead of print].
- Ghisi GLM, Chayes GSDS, Britto RR, Oh P. Health literacy and coronary artery disease: a systematic review. *Patient Educ Couns* 2018;**101**:177–84.
- Miller TA. Health literacy and adherence to medical treatment in chronic and acute illness: a meta-analysis. *Patient Educ Couns* 2016;**99**:1079–86.
- Kannel WB, McGee DL. Diabetes and cardiovascular risk factors: the Framingham study. *Circulation* 1979;**59**:8–13.
- Pujades-Rodriguez M, George J, Shah AD, Rapsomaniki E, Denaxas S, West R, et al. Heterogeneous associations between smoking and a wide range of initial presentations of cardiovascular disease in 1 937 360 people in England: lifetime risks and implications for risk prediction. *Int J Epidemiol* 2015;**44**:129–41.
- Savji N, Rockman CB, Skolnick AH, Guo Y, Adelman MA, Riles T, et al. Association between advanced age and vascular disease in different arterial territories: a population database of over 3.6 million subjects. *J Am Coll Cardiol* 2013;**61**:1736–43.
- Pande RL, Creager MA. Socioeconomic inequality and peripheral artery disease prevalence in US adults. *Circ Cardiovasc Qual Outcomes* 2014;**7**:532–9.
- Cavanaugh KL. Health literacy in diabetes care: explanation, evidence and equipment. *Diabetes Manag (Lond)* 2011;**1**:191–9.
- Husson O, Mols F, Fransen MP, van de Poll-Franse LV, Ezendam NP. Low subjective health literacy is associated with adverse health behaviors and worse health-related quality of life among colorectal cancer survivors: results from the profiles registry. *Psychooncology* 2015;**24**:478–86.
- van der Heide I, Rademakers J, Schipper M, Droomers M, Sørensen K, Uiters E. Health literacy of Dutch adults: a cross sectional survey. *BMC Public Health* 2013;**13**:179.
- Kutner M, Greenberg E, Jin Y, Paulsen C. *The health literacy of America's adults: results from the 2003 national assessment of adult literacy*. Washington, DC: U.S. Department of Education; 2006.
- Mantwill S, Monestel-Umaña S, Schulz PJ. The relationship between health literacy and health disparities: a systematic review. *PLoS One* 2015;**10**:e0145455.
- Weiss BD, Mays MZ, Martz W, Castro KM, DeWalt DA, Pignone MP, et al. Quick assessment of literacy in primary care: the newest vital sign. *Ann Fam Med* 2005;**3**:514–22.
- Fransen MP, Leenaars KE, Rowlands G, Weiss BD, Maat HP, Essink-Bot ML. International application of health literacy measures: adaptation and validation of the newest vital sign in The Netherlands. *Patient Educ Couns* 2014;**97**:403–9.
- Aboumatar HJ, Carson KA, Beach MC, Roter DL, Cooper LA. The impact of health literacy on desire for participation in health-care, medical visit communication, and patient reported outcomes among patients with hypertension. *J Gen Intern Med* 2013;**28**:1469–76.
- Sitzia J, Wood N. Patient satisfaction: a review of issues and concepts. *Soc Sci Med* 1997;**45**:1829–43.
- Spaendonck KPM, Kraaimaat FW. Communicatie in de medische praktijk. *Nijmegen: Afdeling Medische Psychol UMC St Radboud* 2004:15–9.

- 25 Koster ES, Philbert D, Bouvy ML. Health literacy among pharmacy visitors in The Netherlands. *Pharmacoepidemiol Drug Saf* 2015;**24**:716–21.
- 26 Sheridan SL, Halpern DJ, Viera AJ, Berkman ND, Donahue KE, Crotty K. Interventions for individuals with low health literacy: a systematic review. *J Health Commun* 2011;**16**:30–54.
- 27 Peters E, Dieckmann N, Dixon A, Hibbard JH, Mertz CK. Less is more in presenting quality information to consumers. *Med Care Res Rev* 2007;**64**:169–90.
- 28 Volandes AE, Paasche-Orlow MK, Barry MJ, Gillick MR, Minaker KL, Chang Y, et al. Video decision support tool for advance care planning in dementia: randomised controlled trial. *BMJ* 2009;**338**:b2159.
- 29 Kim S, Love F, Quistberg DA, Shea JA. Association of health literacy with self-management behavior in patients with diabetes. *Diabetes Care* 2004;**27**:2980–2.
- 30 Gerber BS, Brodsky IG, Lawless KA, Smolin LI, Arozullah AM, Smith EV, et al. Implementation and evaluation of a low-literacy diabetes education computer multimedia application. *Diabetes Care* 2005;**28**:1574–80.
- 31 Sobel RM, Paasche-Orlow MK, Waite KR, Rittner SS, Wilson EA, Wolf MS. Asthma 1-2-3: a low literacy multimedia tool to educate African American adults about asthma. *J Community Health* 2009;**34**:321–7.
- 32 Wallace AS, Seligman HK, Davis TC, Schillinger D, Arnold CL, Bryant-Shilliday B, et al. Literacy-appropriate educational materials and brief counseling improve diabetes self-management. *Patient Educ Couns* 2009;**75**:328–33.
- 33 Leroy G, Miller T. Perils of providing visual health information overviews for consumers with low health literacy or high stress. *J Am Med Inform Assoc* 2010;**17**:220–3.
- 34 Ross L, Ashford AD, Bleechington SJ, Dark T, Erwin DO. Applicability of a video intervention to increase informed decision making for prostate-specific antigen testing. *J Natl Med Assoc* 2010;**102**:228–36.
- 35 Meade CD, McKinney WP, Barnas GP. Educating patients with limited literacy skills: the effectiveness of printed and videotaped materials about colon cancer. *Am J Public Health* 1994;**84**:119–21.
- 36 Murphy PW, Chesson AL, Walker L, Arnold CL, Chesson LM. Comparing the effectiveness of video and written material for improving knowledge among sleep disorder clinic patients with limited literacy skills. *South Med J* 2000;**93**:297–304.
- 37 Campbell FA, Goldman BD, Boccia ML, Skinner M. The effect of format modifications and reading comprehension on recall of informed consent information by low-income parents: a comparison of print, video and computer-based presentations. *Patient Educ Couns* 2004;**53**:205–16.
- 38 Bryant MD, Schoenberg ED, Johnson TV, Goodman M, Owen-Smith A, Master VA. Multimedia version of a standard medical questionnaire improves patient understanding across all literacy levels. *J Urol* 2009;**182**:1120–5.
- 39 Wilson EA, Park DC, Curtis LM, Cameron KA, Clayman ML, Makoul G, et al. Media and memory: the efficacy of video and print materials for promoting patient education about asthma. *Patient Educ Couns* 2010;**80**:393–8.
- 40 Hironaka LK, Paasche-Orlow MK. The implications of health literacy on patient-provider communication. *Arch Dis Child* 2008;**93**:428–32.
- 41 Meppelink CS, van Weert JC, Haven CJ, Smit EG. The effectiveness of health animations in audiences with different health literacy levels: an experimental study. *J Med Internet Res* 2015;**17**:e11.
- 42 Chew LD, Bradley KA, Boyko EJ. Brief questions to identify patients with inadequate health literacy. *Fam Med* 2004;**36**:588–94.
- 43 Fransen MP, Van Schaik TM, Twickler TB, Essink-Bot ML. Applicability of internationally available health literacy measures in The Netherlands. *J Health Commun* 2011;**16**:134–49.
- 44 Parikh NS, Parker RM, Nurss JR, Baker DW, Williams MV. Shame and health literacy: the unspoken connection. *Patient Educ Couns* 1996;**27**:33–9.